Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 3 February 2008 (MN LATEX style file v2.2)

THE UNIVERSAL ROTATION CURVE OF SPIRAL GALAXIES. II THE DARK MATTER DISTRIBUTION OUT TO THE VIRIAL RADIUS.

P. Salucci[1], A. Lapi[1], C. Tonini[1], G. Gentile[2], I. Yegorova[1], U. Klein[3]
[1] *SISSA International School for Advanced Studies, via Beirut 4, I–34013 Trieste, Italy*
[2] *University of New Mexico, Department of Physics and Astronomy, 800 Yale Blvd NE, Albuquerque, NM 87131, USA*
[3] *Argelander-Institut für Astronomie, Auf dem Hügel 71, D–53121 Bonn, Germany*

3 February 2008

arXiv:astro-ph/0703115v2 8 Mar 2007

ABSTRACT
In the current ΛCDM cosmological scenario, N-body simulations provide us with a Universal mass profile, and consequently a Universal equilibrium circular velocity of the virialized objects, as galaxies. In this paper we obtain, by combining kinematical data of their inner regions with global observational properties, the Universal Rotation Curve (URC) of disk galaxies and the corresponding mass distribution out to their virial radius. This curve extends the results of Paper I, concerning the inner luminous regions of Sb-Im spirals, out to the edge of the galaxy halos.

1 INTRODUCTION

Rotation curves (hereafter RCs) of disk galaxies do not show any Keplerian fall-off and do not match the distribution of the stellar (plus gaseous) matter. As a most natural explanation, this implies an additional invisible mass component (Rubin et al. 1980; Bosma 1981, Persic & Salucci, 1988) that becomes progressively more conspicuous for the less luminous galaxies (e.g.: Persic & Salucci 1988, 1990; Broeils 1992a). Moreover, the kinematical properties of Sb-Im spirals lead to the concept of the *Universal Rotation Curve* (URC) implicit in Rubin 1985, pioneered in Persic and Salucci, 1991 and set in Persic, Salucci & Stel (1996, hereafter PSS, Paper I): RCs can be generally represented out to R_l, the outermost radius where data are available, by $V_{URC}(R;P)$, i.e. by a *universal* function of radius, tuned by some galaxy property P. P can be a global property such as the luminosity and the disk or halo mass or a well defined local quantity like V_{opt}. In any case it serves as the galaxy identifier. In PSS individual RCs and a number of coadded RCs proved the URC paradigm being well fitted by an analytical Curve, $V_{URC}(r/R_{opt}, L)$[1], a function which is the sum in quadrature of two terms: V_{URCD} and V_{URCH}, each representing the disk or halo contribution to the circular velocity:

$$V^2_{URC} = V^2_{URCD} + V^2_{URCH} \tag{1}$$

The stellar component was described by a Freeman disk (Freeman 1970) of surface density $\Sigma_D(r) = \frac{M_D}{2\pi\ R_D^2}\, e^{-r/R_D}$ and contributing to the circular velocity V as:

$$V^2_{URCD}(x) = \frac{1}{2}\frac{GM_D}{R_D}(3.2x)^2(I_0K_0 - I_1K_1) \tag{2a}$$

where $x = r/R_{opt}$ [2] and I_n and K_n are the modified Bessel functions computed at 1.6 x. The dark matter component (with $V^2_{URCH}(r) = \frac{G\,M_H(<r)}{r}$) was described by means of a simple halo velocity profile :

$$V^2_{URCH}(x) = \frac{1}{4\pi} G\rho_0 a^2 x^2/(a^2 + x^2) \tag{2b}$$

The above implies a density profile with an inner flat velocity core of size $\sim aR_{opt}$, a central density ρ_0, an outer r^{-2} decline. The sum of the contributions (2a) and (2b) well fit all the PSS data with ρ_0, a^2 specific functions of luminosity (see PSS). Let us remind that disk masses M_D of spirals were found in the range $10^9\ \mathrm{M}_\odot \leq M_D \leq 2\times 10^{11}\ \mathrm{M}_\odot$.

[1] The reader is directed to PSS for the details of the procedure.
[2] We define the "disk size" $R_{opt} \equiv 3.2\, R_D$

The URC for the purpose of this work matches well the individual RCs of late type spirals (see also Appendix for a discussion). It is useful to express the URC paradigm in the following way: at any chosen radius, the URC predicts the circular velocity of a (late type) spiral of *known* luminosity and disk scale-length, within an error that is one order of magnitude smaller than the variations it shows i) at different radii and ii) at any radius, with respect to objects of different luminosity.

Let us remind that the Universal curve built in PSS holds out to R_l, uses the luminosity as the galaxy identifier and the disk scale-length as a unit of measure for the radial coordinate. We will label it as URC_0 to indicate it as the first step of a definitive function of the dark radial coordinate, able to reproduce the observed RCs of spirals. URC_0 provides fundamental knowledge on the mass distribution in spirals, while it suffers from three limitations: 1) it strictly holds only in a region extended less than 5% the DM halo size (see below) 2) the velocity profile of the halo component, valid out to R_l, cannot be extrapolated to radii of cosmological interest 3) it identifies objects by their luminosities, rather than by their virial masses. Let us point out that the URC_0 has been often and successfully used as an observational benchmark for theories, but this, only for $R < R_l$ and after that a relation between the halo mass and the galaxy luminosity was assumed.

On the other side, high–resolution cosmological N–body simulations have shown that, within the (Λ) Cold Dark Matter (CDM) scenario, dark halos achieve a specific equilibrium density profile characterized by a universal shape and, in turn, an universal halo circular velocity (Navarro, Frenk & White, 1997, NFW), $V_{\rm NFW}(R, M_{vir})$ in which the virial mass M_{vir} and virial radius R_{vir} are the galaxy identifier and radial coordinate.

$$\rho_H(r) = \frac{M_{vir}}{4\pi R_{vir}^3}\,\frac{c^2\,g(c)}{x\,(1+cx)^2}\ , \qquad (3a)$$

where $x \equiv r/R_{vir}$ is the radial coordinate, c is the concentration parameter, and $g(c) = [\ln(1+c) - c/(1+c)]^{-1}$. The parameter c is found to be a weak function of the halo mass, given by $c \approx 14\,\left(M_{vir}/10^{11}\ \mathrm{M}_\odot\right)^{-0.13}$ (Bullock et al, 2001, Dutton, 2006, Gnedin 2006). This leads to

$$V^2_{\rm NFW}(r) = V^2_{vir}\frac{c}{g(c)}\frac{g(x)}{x}\ , \qquad (3b)$$

with $V_{vir} = V(R_{vir})$. It is interesting to note that in this scenario the present–day circular velocity, which also includes a baryonic component arranged in a disk, is predicted to be a *Universal* function of radius, tuned by few galaxy parameters (Mo, Mao & White 1998). However, it is well known that observations of spiral galaxies favor density concentrations lower than those predicted for CDM by Eq. (3a): DM halos detected around spirals do not show the NFW central cusp in favor of a core-like structure (Gentile et 2005; van den Bosch & Swaters 2001; Swaters et al. 2003; Weldrake et al. 2003; Simon et al. 2005; Donato et al, 2004; Gentile et al. 2007).

Therefore, the reconstruction of the mass distribution of DM halos from observations *in parallel* with that emerging from N-body simulations is required not only as a normal scientific routine, but also in view of a theory-vs-observations likely disagreement.

As an alternative to the simulation method, we will support the URC paradigm by means of a set of proper observational data and we will derive an analytical form for this curve, valid from the galaxy center out to its virial radius and characterized by the halo mass as the galaxy identifier. In detail, we extend/improve the URC_0 in PSS a) by adopting a different halo profile, proper to describe the halo distribution out to the virial radius, b) by using a number of RCs substantially more extended than those in PSS and c) by exploiting the relationship between the disk mass M_D, and the virial galaxy mass M_{vir}, recently obtained by Shankar et al. (2006). This will allow to build an "observational" Universal Curve, $V_{URC}(R; M_{vir})$, extended out to R_{vir} and having the virial mass as the galaxy identifier. This curve is the observational counterpart of the universal ΛCDM NFW N-body generated profile.

While pointing that the concept behind the Universal Rotation Curve may be valid also for galaxies of different Hubble Types (see Salucci and Persic, 1997), but a number of issues are still open and will be dealt elsewhere:

i) Sa galaxies amount, by number, to less than 10% of the whole spiral population, and are important objects in view of the dual nature of their stellar distribution. They show RC profiles with a clear systematics with luminosity (Rubin et al. , 1985), but, not unexpectedly, with some difference from those of the URC_0 (Noordmeer, 2007).

ii) Dwarf spirals with $V_{opt} < 50 km/s$ are not well studied and included in the URC yet, also because in these objects the RCs do not coincide with the circular velocity, being significant the complex asymmetric drift correction.

iii) The kinematical properties of spirals of very high stellar disk mass are not presently investigated with a suitably large sample.

iv) A possible additional URC physical parameter (e.g. the surface stellar density) to take care of the (small) variance of the RCs profiles that seems to be unaccounted by the luminosity.

Finally, let us remind that, in a flat cosmology with matter density parameter $\Omega_M = 0.27$ and Hubble constant $H_0 = 71$ km s^{-1} Mpc^{-1}, at the present time, the halo virial radius R_{vir}, i.e. the size of the virialized cosmological perturbation of mass M_{vir} scales with the latter as:

$$R_{vir} = 259\Big(\frac{M_{vir}}{10^{12}\ \mathrm{M}_\odot}\Big)^{1/3}\ \mathrm{kpc}\ . \qquad (4)$$



Figure 1. The core radius vs. virial mass relations for the SE sample and the present work (solid line).

(see e.g. Eke et al. 1996).

2 THE UNIVERSAL HALO VELOCITY PROFILE

We assume that the DM mass distribution is described by the Burkert (1995) profile

$$\rho(r) = \frac{\rho_0\, r_0^3}{(r + r_0)\,(r^2 + r_0^2)}\ , \tag{5a}$$

where r_0 is the core radius and ρ_0 the effective core density, in principle two independent parameters. Correspondingly, the total halo mass inside radius r is given by $M_H(r) = 4\,M_0\,\left[\ln\left(1+\frac{r}{r_0}\right) - \tan^{-1}\left(\frac{r}{r_0}\right) + \frac{1}{2}\,\ln\left(1+\frac{r^2}{r_0^2}\right)\right]$ with $M_0 = 1.6\,\rho_0\,r_0^3$, so that:

$$V^2_{URCH}(r) = 6.4\ G\ \frac{\rho_0 r_0^3}{r}\left\{ln\left(1+\frac{r}{r_0}\right) - \tan^{-1}\left(\frac{r}{r_0}\right) + \frac{1}{2}ln\left[1+\left(\frac{r}{r_0}\right)^2\right]\right\}\ . \tag{5b}$$

Inside R_l this profile is indistinguishable from the halo term (2b) in the URC$_0$ (Salucci & Burkert, 2000, Gentile et al 2004). At larger radii, the mass diverges only logarithmically with radius and converges to the NFW velocity profile, provided that $r_0 \ll R_{vir}$.

We fit the set of individual and coadded RCs of PSS with $V_{URC}(R; M_D, \rho_0, r_0)$ and derive the model parameters M_D, ρ_0, r_0 (see Salucci and Burkert, 2001):

$$\log \frac{\rho_0}{\text{g cm}^{-3}} = -23.515 - 0.964\,\left(\frac{M_D}{10^{11}\,\text{M}_\odot}\right)^{0.31} \tag{6a}$$

and

$$\rho_0 = 5\times 10^{-24} r_0^{-2/3} e^{-(r_0/27)^2}\,\text{g cm}^{-3}\ . \tag{6b}$$

Eqs. (1)-(2a)-(5b)-(6) define the URC out to R_l, $V_{URC}(R, M_D, r_0)$, from the "baryonic perspective". Let us notice that, as result of the RC mass modeling, r_0, differently from M_D and ρ_0, has quite large fitting uncertainties, viz. $\delta r_0/r_0 = 0.3 - 0.5$. Following our empirical approach, we do not extrapolate the URCH determined inside R_l out to $R_{vir} \gg R_l$. In that this will be uncertain besides of unknown validity. This quantity will be derived in the next section.

3 THE URC OUT TO THE VIRIAL RADIUS

We overcome the two main limitations of the URC$_0$, its problematic extrapolation between R_l and R_{vir} and the uncertainty in the estimate of the core radius, by determining the latter by means of a new *outer* observational quantity, the halo virial velocity $V_{vir} \equiv [GM_{vir}/R_{vir}(M_{vir})]^{1/2}$, related to the virial mass through Eq. (4). In detail, we obtain V_{vir} from the disk mass, suitably measured from inner kinematics through its relationship to the virial mass found by Shankar et al. (2006)

that is well represented (i.e. within 5%, the actual relation we use is shown in Fig. 2 and given in the Code indicated in the Discussion) by:

$$M_D = 2.3 \times 10^{10}\ M_\odot \frac{[M_{vir}/(3\ 10^{11}\ \mathrm{M}_\odot)]^{3.1}}{1+[M_{vir}/(3\ 10^{11}\mathrm{M}_\odot)]^{2.2}}\ . \tag{7}$$

This relationship is a consequence of the existence of i) the universal stellar mass function, (Bauldry et al 2004, Bell et al 2003) and of ii) the cosmological halo mass function as indicated by N-body ΛCDM simulations .[3] Let us notice that this relationship is obtained without assuming any halo density profile, so that it can be combined with the mass modelling of the inner kinematics.

Let us first derive $R_D(M_{vir})$, the disk scale-length as a function of the halo mass, by inserting Eq. (7) in the relationship:

$$\log \frac{R_D}{\mathrm{kpc}} = 0.633 + 0.379\ \log \frac{M_D}{10^{11}\ \mathrm{M}_\odot} + 0.069\ \left(\log \frac{M_D}{10^{11}\ \mathrm{M}_\odot}\right)^2 \tag{8}$$

obtained in PSS. We note that no result of this work is affected by the observational uncertainties on the relationship Eq. (8).

It is worth to compute the radial extrapolation needed to reach R_{vir} from $R_l = 6R_D$, a quantity that can also be associated with the baryonic collapse factor $F = R_{vir}/R_D$; we find $F \approx 90 - 15\ log\frac{M_{vir}}{10^{11}M_\odot}$, i.e. about 25 times the disk size $\sim 3R_D$.

Eq. (7) in combination with Eq. (4) allows us to add, to the PSS set of kinematical data leading to the URC$_0$, a new observational quantity: $V_{vir}(M_D) = GM_{vir}/R_{vir}$, relative to the virial radius. Then, we determine the core radius not from the inner kinematics, but as the value of r_0 for which the velocity model described by Eqs. (1), (2a), (5b) and (6a), matches (at R_{vir}) the virial velocity V_{vir} given by Eq. (7) and (4). Let us write this as:

$$\frac{GM_{vir}}{R_{vir}(M_{vir})} = V^2_{URCH}[R_{vir}(M_{vir}); \rho_0(M_{vir}), r_0)] \tag{9a}$$

where $\rho(M_{vir})$ as a short form for $\rho(M_D(M_{vir}))$ with Eq. (7) inserted in Eq. (6a). From the PSS inner kinematics we get the values of ρ_0 and M_D according to Eq. (6a), so that Eq. (9a) becomes an implicit relation between r_0 and M_{vir} (c_1, c_2 are known numerical constants):

$$M_{vir} = c_1 \frac{\rho_0(M_{vir}) r_0^3}{c_2 M_{vir}^{1/3}} \left\{ ln\left(1+\frac{c_2 M_{vir}^{1/3}}{r_0}\right) - \tan^{-1}\left(\frac{c_2 M_{vir}^{1/3}}{r_0}\right) + \frac{1}{2} ln\left[1+\left(\frac{c_2 M_{vir}^{1/3}}{r_0}\right)^2\right]\right\}\ . \tag{9b}$$

The above can be numerically solved for any M_{vir}, and the solution can be approximated by:

$$\log\left(r_0/\mathrm{kpc}\right) \simeq 0.66 + 0.58\ \log\left(M_{vir}/10^{11}\ \mathrm{M}_\odot\right)\ . \tag{10}$$

(an higher order approximation is given in the Code indicated in the Discussion). Let us stress that the present derivation of r_0 is very solid with respect to observational uncertainties: errors up to a factor 2 in M_{vir} in eq (7) trigger errors in r_0 lower than 40%, and errors in the outer halo velocity slope ($0.1 \leq R/R_{vir} \leq 1$) lower than 0.1. This is certainly smaller than the scatter of values with which this quantity is found by N-Body simulations and by SPH/semi analytical studies of galaxy formation including the baryonic components.

It is worth investigating a number of recently published super-extended (SE) RCs (Donato et al. 2005, Gentile et al 2004, Salucci et al 2003). They reach a radius larger than 5% (and up to 15%) of the virial radius, i.e. a radius at least twice as extended as those of the sinthetic curves in PSS. The mass modelling of these SE RCs (made in the original papers) shows an r_0 vs. M_{vir} relationship that is in good agreement with Eq. (10). Relation (10) and the above individual values differ by 20% -40% from those determined from the inner kinematics alone and given by means of eq (6b). Since in this paper (also because $R_l \sim r_0$), eq (6) is considered a prediction of the inner mass modelling rather than an actual measurement, such good agreement indicates the soundness of the PSS mass modelling.

Let us notice that only for a range of values of the crucial quantity $V(R_l)$- V_{vir}, with the first term obtained by the inner kinematics and the second one via Eqs. (4) and (7), there is a solution for eq.(9b), therefore, the existence of eq (10) and the agreement of the values of the Burkert core radii, measured independently at $0.05R_{vir}$, $0.1R_{vir}$ and R_{vir} are important tests passed by this profile.

Then, by means of Eqs. (1), (2a)-(5b)-(6a)-(10), we construct the full URC, extended out to the virial radius and with the virial mass as the galaxy indicator. It is useful to show the relationships we use (see Fig. 2). The mass model includes a Burkert DM halo of central density ρ_0, of core radius of size r_0 and a Freeman disk of mass M_D. The URC fits nicely the available velocity data out to R_l and it is valid out to the virial radius, where it exactly matches V_{vir}. Moreover, since M_{vir} is the quantity that in theoretical studies identifies a galaxy, we overcome the main limitation of URC$_0$.

We consider all of the three coordinate systems $r, r/R_D, r/R_{vir}$ equivalent to represent the main structural properties of

[3] There is no inconsistency in adopting the ΛCDM halo mass function and cored halo mass models, in that the latter can be formed astrophysical from the cosmological cuspy ones. Since both functions account the same cosmological objects, the Jacobian of their transformation defines a relation between the disk and virial mass in spirals (see Shankar et al. 2006).



Figure 2. The various relationships used in this paper.



Figure 3. The Universal Rotation Curve in physical units. Each curve corresponds to $M_{vir} = 10^{11}\ 10^{n/5} \mathrm{M}_\odot$, with $n = 1 \ldots 9$ from the lowest to the highest curve.

the mass distribution in spirals, but each of them showing some particular aspects. More specifically, it is then possible and useful to build several "URCs" i.e. $V_{URC}(r/R_{coo}; P)$, where P is a galaxy identifier (M_D, M_{vir}, L) and R_{coo} a radial coordinate $(r, r/R_D, r/R_{vir})$. Although not all these URCs are independent in a statistical sense, they all are relevant in that they all well reproduce the individual RCs and each of them highlights particular properties of the mass distribution.

In Fig. 3 we show $V_{URC}(r; M_{vir})$, the URC in physical units with the objects identified by the halo virial mass; each line refers to a given halo mass in the range $10^{11}\,\mathrm{M}_\odot \lesssim \mathrm{M_{vir}} \lesssim 10^{13}\,\mathrm{M}_\odot$; the halo mass determines both the amplitude and the shape of the curve. Note the contribution of the baryonic component, negligible for small masses but increasingly important in the larger structures, mirrors the behavior of the $M_{vir} - M_D$ relation. The general existence of an inner peak is evident but, especially at low masses, it is due to both dark and stellar components. Remarkably, the maximum value of the circular velocity occurs at about 15 ± 3 kpc, independent of the galaxy mass: this seems to be a main kinematic imprint of the DM - luminous mass interaction occurring in spirals. Furthermore, Fig. 3 shows that the "Cosmic Conspiracy" paradigm has no observational support: there is no fine tuning between the dark and the stellar structural parameters to produce the same particular RC profile in all objects (e.g. a flat one). Conversely, a number of relationships between the various structural parameters produce



Figure 4. The Universal Rotation Curve, normalized at its virial value $V_{URC}(R_{vir})$, as a function of normalized dark radius $x \equiv R/R_{vir}$. Each curve, from the highest to the lowest, corresponds to M_{vir} defined as in Fig. 3. The bold line is the NFW velocity profile (see text).

a variety of RC profiles. Moreover, the peak velocity of the stellar component $V_{disk}^{peak} = V_D(2.2R_D) = GM_D/R_D\ k$, with $k = const$, is not a constant fraction of the virial velocity as is found in ellipticals, (i.e $\sigma \propto V_{vir}$), but it ranges between the values 1 and 2 depending on the halo mass.

Moreover, as in the NFW (and Burkert) RC profiles, the URC profiles are found (moderately) decreasing over most of the halo radial extent. The paradigm of flat rotation curves is obviously incorrect even/especially intended as an asymptotic behavior at large radii. In fact, we find that both $V(0.05R_{vir})$,the velocity at the farthest radius with available kinematical in PSS and $V(3R_D)$, a main reference velocity of the luminous regions of spirals, are significantly $(10\% - 30\%)$ higher than the (observational) value of V_{vir}. This rules out a $V = constant$ extrapolation of the inner RCs out to regions non mapped by the kinematics and DM dominated regions. We note that this result is independent of the adopted halo density profile and is far from being granted on theoretical grounds.

In Fig. 4 we frame the URC from a full DM perspective by plotting $V_{URC}(R/R_{vir}; M_{vir})$. We set the virial mass M_{vir} as the galaxy identifier and R/R_{vir} as the radial "dark" coordinate, thus normalizing the amplitudes by $V_{vir} \propto M_{vir}^{1/3}$. This ensemble of curves, a main goal of the present work, is parallel to those emerging in N-body simulations and aims to represent the actual velocity profiles of spirals. In these variables the DM halos are self-similar; the whole system is self-similar in the outer regions, while in the innermost 30% of the halo size the baryons have influenced the dynamics and broken the self-similarity. In these coordinates it easily emerges that the maximum of the RC occurs at very different radii, viz. at $\simeq 2R_D$ for the most massive objects and at $\sim 10R_D$ for the least massive ones. Then, no reference circular velocities, to be considered as the actual physical counterparts of the empirical velocities of the Tully-Fisher relationship, exist in actual galaxy rotation curves.

In Fig. 5 we zoom into the URC to look for the inner (luminous) regions of spirals from a baryonic perspective: the URC is so expressed as a function of the "baryonic" radial coordinate r/R_D. This figure corresponds to Fig. 4 of PSS, with the important difference that here the virial mass, rather than the galaxy luminosity, is the galaxy identifier. Plainly, an inverse correlation between the average steepness of the RC slope and the halo mass holds, similar to the slope-luminosity relationship found by Persic & Salucci (1988). In this coordinate the stellar matter is closely self-similar, and the different shapes of the RC's curves are mainly due to the $M_{vir} - M_D$ relation. In the space defined by normalized circular velocity - dark radius - halo mass, spirals do not occupy random positions, but a well defined plane of very small thickness. We clearly see that, by filling only less than $10^{-3}\%$ of the available volume, the available kinematics of spiral galaxies defines the Universal Rotation Curve. Let us notice that, in principle, theories of the formation of spirals do not trivially imply the existence of such a surface that underlies the occurrence of a strong dark - luminous coupling.

We now show the URC DM density distribution. In Fig. 6 we show it as a function of x and M_{vir}. For $x < 10^{-1}$ the well known core-cusp discrepancy emerges, i.e. the DM density of actual halos around spirals is about one order of magnitude smaller and radially much more constant than the NFW predictions. At $x > 0.4$, for a concentration parameter $c = 13(M_{vir}/10^{12}\ M_\odot)^{-0.13}$, the observed halo densities are consistent with the NFW predictions for halos of the same virial mass. Note that this is a direct test: for halos with density profiles at $x > 0.5$ very different from the Burkert or the NFW profiles, Eq. 9b does not have solution.

More specifically, let us constrain the analytical form of the outer DM distribution. For $2 \leq r/r_0 \leq 18$, the following



Figure 5. The inner Universal Rotation Curve, normalized at its value at $4R_D$, as a function of normalized stellar radius R/R_D for galaxies with M_{vir} as in Fig. 3.



Figure 6. The URC halo density vs. the NFW halo density of objects of the same mass, as a function of normalized radius and virial mass. The axes labels are x, $logM_{vir}/M_{\odot}$ and $log(\rho/(g\ cm^{-3}))$

approximation for the Burkert and NFW profile holds ($y \equiv r/r_0$, $\epsilon = 0$)

$$V_{URCH}(y) = V_{URCH}(3.24)\frac{2.06\, y^{0.86}}{1.59 + y^{1.19+\epsilon}}\ . \tag{11}$$

Let us suppose that the actual *outer* DM velocity *profile* is different from the Burkert/NFW given by Eq. (11), i.e. $\epsilon \neq 0$. Then in Fig 7 we show that, even assuming large uncertainties in V_{vir}, in order to match both $V(R_l)$ and V_{vir}, we must have $\epsilon < 0.1$. This is a first direct support for the Burkert and the NFW density law to be able to represent the outer regions ($0.3 \leq r/R_{vir} \leq 1$) of DM galaxy halos.

Notice that weak-lensing shear fields, at several hundreds kpc from the galaxy centers, are found *compatible* with the predictions of the NFW density profile, but cannot exclude non-NFW profiles (Kleinheinrich ., et al., 2006 and references therein).

4 DISCUSSION AND CONCLUSIONS

In this paper, we have built the Universal Rotation Curve of spiral galaxies by means of kinematical and photometric data. We physically extended the URC, established for the inner region of galaxies in PSS out to R_{vir} and have been able to employ the virial mass M_{vir} as the parameter that characterizes spiral galaxies, and the virial radius R_{vir} as a unit of measure for the radial coordinate. This URC is meant to be the *observational* counterpart of the NFW rotation curve, emerging from cosmological simulations performed in the CDM scenario. The URC yields the gravitational potential at any radius and it allows to link the local properties in the inner luminous regions with the global properties of the DM halos.

DM halos have one (and likely just one) characteristic length scale, $r_0 \propto M_{vir}^{0.6}$, which it is not *naturally* present in current scenarios of galaxy formation. Thus, they do not show any sign of an inner cuspy region of size $r_s \propto M_{vir}^{0.4}$. The halo velocity contribution V_{URCH} rises with radius like a solid body at $r \sim 0$, decelerates to reach a maximum at 3.24 r_0 from where it start to slowly decrease out to R_{vir} with a slope that it is consistent with that of the NFW halos. The main significance of the URC concerns the full mass distribution (MD). First, it is possible to immediately exclude the following scenarios (and combinations of them): i) individual behavior, every object has its own MD; ii) unique behavior, every object has almost the same MD. Instead, the MD in spirals shows a remarkable *mass-dependent* systematics: both the dark and the stellar matter are distributed according to *profiles* that are functions of the total mass M_{vir} (see Fig. 8). Finally, the DM halo becomes the dominant mass component in galaxies at different radii, according to the galaxy mass: from $\sim 10^{-2} R_{vir}$ for the lowest masses, to $\sim 10^{-1} R_{vir}$ for the highest ones.

We write the equilibrium velocity of the halos around spirals as the following approximation of the relations in the previous Sect.:

$$V_{URCH} = A(M_{vir}) x^{-1/2} \Big\{ ln\Big[1 + \gamma(M_{vir})x\Big] - \tan^{-1}\Big[\gamma(M_{vir})x\Big] + \frac{1}{2} ln\Big[1 + \gamma(M_{vir})x^2\Big] \Big\}^{0.5}$$

with $A(M_{vir}) = 0.406 + 1.08\ log[M_{vir}/(10^{11}\mathrm{M}_\odot)] - 0.688\ \{\log[\mathrm{M_{vir}}/(10^{11}\mathrm{M}_\odot)]\}^2) + 0.766\ \{\log[\mathrm{M_{vir}}/(10^{11}\mathrm{M}_\odot)]\}^3$ and $\gamma(M_{vir}) = 26.78\ [M_{vir}/(10^{11}\mathrm{M}_\odot)]^{-0.246}$. This is the observational counterparts of N-body outcomes.

A Mathematica code for the figures in this paper is available at: http://www.novicosmo.org/salucci.asp.

5 APPENDIX

In this Appendix we discuss the observational evidence for the URC claim, the nature and the implications of which it is worth to clarify. The paradigm states that, when binned by luminosity, the RCs form a set of smooth, low-scatter synthetic curves, whose profiles and amplitudes are strong functions of the luminosity bin. [4]. Furthermore, the URC paradigm implies: i) rotation velocity slopes *vs.* rotation velocity amplitudes relationships (see fig (2) and (3) of PSS) and ii) a set of relations (Radial Tully Fisher relationship) holding at different radii x, defined as:

$$log\ V(x) = a_x M + b_x$$

where $x \equiv R/R_D$ and a_x and b_x are the fitting parameters, and M is the galaxy magnitude (Persic and Salucci 1991).

Evidence for the URC claim and/or its above implications comes from: a) detailed analyzes of independent samples: Catinella et al 2006, (2200 RC's, see their Fig. 12), Swaters 1999, (60 extended RC's, see Chapter 4); b) independent analyzes of the PSS sample: Rhee (1996), Roscoe (1999); c) the finding of a very tight RTF in PSS and other three different samples Willick (1997, see below), Yegorova et al. (2007).

The claim has been also tested by comparing the RCs of two samples of spirals (Courteau, 1998, 131 objects; Verheijen,

[4] The *analytical form* of the URC is built by assuming reasonable disk-halo velocity profiles, with three free parameters $(V(R_{opt}), a, \beta)$ that are obtained by χ^2-fitting the sinthetic curves.



Figure 7. Halo velocities at R_l and R_{vir} (*filled circles*) vs. the URC-halo velocity, given by Eqs. (11) (*solid line*) and vs. velocity profiles with average logarithmic slope steeper or shallower by an amount $\epsilon = 0.1$ (*dashed line*).



Figure 8. The dark halo (solid line), the disk (short dashed line), and the total (long dashed) URC mass profile for reference masses of $10^{11}M_\odot$ and $10^{13}M_\odot$. The distribution for the intermediate halo masses can be derived from section 2

1997, 30 objects) with the circular velocities predicted by the URC_0, once that the values of galaxy luminosity and disk length-scale are inserted in it. The face-value result of the test: 2/3 of the RCs are in pretty good agreement with the universal curve, while 1/3 show some disagreement, indicates that the URC is a useful tool to investigate the systematics of the mass distribution in spirals, but also it questions about its universality. However, while some of this disagreement may reflect an inefficiency of the URC_0 to reproduce the RCs, the actual performance of the URC is better than it is claimed. In fact, spurious data vs predictions disagreements are created in performing this test and precisely when they insert in URC_0 the

values of L_B and R_D, affected by (occasionally large) observational errors. By taking into account this effect the URC_0 success rate reaches 80% and more.

Willick (1998) found, by studying a large sample of RCs, a radial variation of the scatter of the inverse RTF defined above and he interpreted it as an evidence against the URC. Let us show that this argument is incorrect and that *au contraire* the properties of the RTF support the URC paradigm. The increase/decrease of the scatter found is very small (Willick, 1998): the scatter ranges from 0.065 dex (at $2R_D$) to 0.080 dex (at $0.5R_D$ and at $3R_D$) and it implies, if totally intrinsic, a prediction error in $log\ V(x)$ of $(0.08^2 - 0.065^2)^{0.5} = 0.04\ dex$. Moreover, some of the scatter increase/decrease is due to the larger random observational errors present in the outermost measurements; in fact, a refined analysis of the issue (Yegorova et al 2007) finds a smaller predicting error for three large sample of spirals. Therefore, from the RTF we have that, in the region considered, the luminosity *statistically* predicts the circular velocity at any radius and in any galaxy within an error of 5% - 10% , a quantity much smaller than the variations of the latter *in each* galaxy and *among* galaxies.

ACKNOWLEDGEMENTS

We thank L. Danese for helpful discussions, and the referee for useful comments.

6 REFERENCES

Baldry, I.K., et al. 2004, ApJ, 600, 681
Bell, E.F., McIntosh, D.H., Katz, N., & Weinberg, M.D. 2003, ApJS, 149, 289
Borriello, A., Salucci, P. 2001, MNRAS, 323, 285
Bosma, A. 1981, AJ, 86, 1791
Broeils, A.H. 1992a, A&A, 256, 19
Broeils, A.H. 1992b, Ph.D. thesis, Groningen University
Catinella, B., Giovanelli R., Haynes, M.P., 2006, ApJ. 640, 751
Courteau S., 1997, AJ, 114, 2402
Donato, F., Gentile, G., Salucci, P., 2004, MNRAS, 353, 17
Eke, V.R., Cole, S., & Frenk, C.S., 1996, MNRAS 282,263
Freeman, K.C. 1970, ApJ, 160, 811
Gentile, G., Burkert, A., Salucci, P., Klein, U., Walter, F. 2005, ApJ, 634, L145
Gentile G., Salucci P., Klein U., Vergani D., Kalberla P., 2004, MNRAS, 351, 903
Gentile G., Salucci P., Klein U., Granato G.L., 2007, MNRAS, 375, 199
Kleinheinrich M., et al., 2006, A&A, 455,441
Mo, H.J., Mao, S., White, S.D.M. 1998, MNRAS, 295, 319
Noordermeer, E, 2007, MNRAS, in press, astro-ph/0701731
Navarro, J.F., Frenk, C.S., & White, S.D.M. 1996, ApJ, 462, 563
Persic, M., Salucci, P. 1988, MNRAS, 234, 131 (PS88)
Persic M., Salucci P., 1990, MNRAS, 245, 577
Persic, M., Salucci, P. 1991, ApJ, 368, 60 (PS91)
Persic, M., Salucci, P. 1995, ApJS, 99, 501 (PS95)
Persic, M., Salucci, P., Stel, F. 1996, MNRAS, 281, 27
Rhee, M-H 1997, Thesis, Groningen University.
Roscoe, D. F. 1999, A&A **343**, 788
Rubin, V.C., Ford, W.K., Jr., Thonnard, N. 1980, ApJ, 238, 471
Rubin, V.C., Ford, W.K., Jr., Thonnard, N., Burstein, D., 1982, ApJ, 261, 439
Rubin V. C., Burstein D., Ford W.K., Jr., Thonnard N., 1985, ApJ, 289, 81
Salucci P., Burkert A., 2000, ApJ, 537, L9
Salucci P., Persic M., 1997, ASPC, 117, 1
Shankar, F., Lapi, A., Salucci, P., De Zotti, G., Danese, L. 2006, ApJ, 643, 14
Simon, J. D., Bolatto, A. D., Leroy, A., Blitz, L., Gates, E. L. 2005, ApJ, 621, 757
Swaters, R.A., Madore, B.F., van den Bosch, F.C., Balcells, M. 2003, ApJ, 583, 732
Swaters, 1999, Ph.D. Thesis, Groningen University
Tonini, C., Lapi, A. & Salucci, P. 2006, ApJ, in press
Tonini, C., Lapi, A., Shankar, F. & Salucci, P. 2006, ApJ, 638, L13
van den Bosch, F.C., Swaters, R.A., 2001 MNRAS, 325, 1017
Verheijen, M, 1997, Ph.D. Thesis, Groningen University

Yegorova, I. et al, 2007, MNRAS in press, astro-ph/0612434
Weldrake, D.T.F., de Blok, W. J. G., Walter, F. 2003, MNRAS, 340, 12
Willick J.A., 1999, ApJ, 516, 47

Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 3 February 2008 (MN LATEX style file v2.2)

THE UNIVERSAL ROTATION CURVE OF SPIRAL GALAXIES: II THE DARK MATTER DISTRIBUTION OUT TO THE VIRIAL RADIUS.

P. Salucci[1], A. Lapi[1], C. Tonini[1], G. Gentile[2], I. Yegorova[1], U. Klein[3]
[1] *SISSA International School for Advanced Studies, via Beirut 4, I–34013 Trieste, Italy*
[2] *University of New Mexico, Department of Physics and Astronomy, 800 Yale Blvd NE, Albuquerque, NM 87131, USA*
[3] *Argelander-Institut für Astronomie, Auf dem Hügel 71, D–53121 Bonn, Germany*

3 February 2008

arXiv:astro-ph/0703115v2 8 Mar 2007

ABSTRACT
In the current ΛCDM cosmological scenario, N-body simulations provide us with a Universal mass profile, and consequently a Universal equilibrium circular velocity of the virialized objects, as galaxies. In this paper we obtain, by combining kinematical data of their inner regions with global observational properties, the Universal Rotation Curve (URC) of disk galaxies and the corresponding mass distribution out to their virial radius. This curve extends the results of Paper I, concerning the inner luminous regions of Sb-Im spirals, out to the edge of the galaxy halos.

1 INTRODUCTION

Rotation curves (hereafter RCs) of disk galaxies do not show any Keplerian fall-off and do not match the distribution of the stellar (plus gaseous) matter. As a most natural explanation, this implies an additional invisible mass component (Rubin et al. 1980; Bosma 1981, Persic & Salucci, 1988) that becomes progressively more conspicuous for the less luminous galaxies (e.g.: Persic & Salucci 1988, 1990; Broeils 1992a). Moreover, the kinematical properties of Sb-Im spirals lead to the concept of the *Universal Rotation Curve* (URC) implicit in Rubin 1985, pioneered in Persic and Salucci, 1991 and set in Persic, Salucci & Stel (1996, hereafter PSS, Paper I): RCs can be generally represented out to R_l, the outermost radius where data are available, by $V_{URC}(R;P)$, i.e. by a *universal* function of radius, tuned by some galaxy property P. P can be a global property such as the luminosity and the disk or halo mass or a well defined local quantity like V_{opt}. In any case it serves as the galaxy identifier. In PSS individual RCs and a number of coadded RCs proved the URC paradigm being well fitted by an analytical Curve, $V_{URC}(r/R_{opt},L)$[1], a function which is the sum in quadrature of two terms: V_{URCD} and V_{URCH}, each representing the disk or halo contribution to the circular velocity:

$$V^2_{URC} = V^2_{URCD} + V^2_{URCH} \tag{1}$$

The stellar component was described by a Freeman disk (Freeman 1970) of surface density $\Sigma_D(r) = \frac{M_D}{2\pi\ R_D^2}\, e^{-r/R_D}$ and contributing to the circular velocity V as:

$$V^2_{URCD}(x) = \frac{1}{2}\frac{GM_D}{R_D}(3.2x)^2(I_0K_0 - I_1K_1) \tag{2a}$$

where $x = r/R_{opt}$ [2] and I_n and K_n are the modified Bessel functions computed at 1.6 x. The dark matter component (with $V^2_{URCH}(r) = \frac{G\,M_H(<r)}{r}$) was described by means of a simple halo velocity profile :

$$V^2_{URCH}(x) = \frac{1}{4\pi}G\rho_0 a^2x^2/(a^2+x^2) \tag{2b}$$

The above implies a density profile with an inner flat velocity core of size $\sim aR_{opt}$, a central density ρ_0, an outer r^{-2} decline. The sum of the contributions (2a) and (2b) well fit all the PSS data with ρ_0, a^2 specific functions of luminosity (see PSS). Let us remind that disk masses M_D of spirals were found in the range $10^9\ \mathrm{M}_\odot \leq M_D \leq 2\times10^{11}\ \mathrm{M}_\odot$.

[1] The reader is directed to PSS for the details of the procedure.
[2] We define the "disk size" $R_{opt} \equiv 3.2\,R_D$

The URC for the purpose of this work matches well the individual RCs of late type spirals (see also Appendix for a discussion). It is useful to express the URC paradigm in the following way: at any chosen radius, the URC predicts the circular velocity of a (late type) spiral of *known* luminosity and disk scale-length, within an error that is one order of magnitude smaller than the variations it shows i) at different radii and ii) at any radius, with respect to objects of different luminosity.

Let us remind that the Universal curve built in PSS holds out to R_l, uses the luminosity as the galaxy identifier and the disk scale-length as a unit of measure for the radial coordinate. We will label it as URC_0 to indicate it as the first step of a definitive function of the dark radial coordinate, able to reproduce the observed RCs of spirals. URC_0 provides fundamental knowledge on the mass distribution in spirals, while it suffers from three limitations: 1) it strictly holds only in a region extended less than 5% the DM halo size (see below) 2) the velocity profile of the halo component, valid out to R_l, cannot be extrapolated to radii of cosmological interest 3) it identifies objects by their luminosities, rather than by their virial masses. Let us point out that the URC_0 has been often and successfully used as an observational benchmark for theories, but this, only for $R < R_l$ and after that a relation between the halo mass and the galaxy luminosity was assumed.

On the other side, high–resolution cosmological N–body simulations have shown that, within the (Λ) Cold Dark Matter (CDM) scenario, dark halos achieve a specific equilibrium density profile characterized by a universal shape and, in turn, an universal halo circular velocity (Navarro, Frenk & White, 1997, NFW), $V_{\rm NFW}(R, M_{vir})$ in which the virial mass M_{vir} and virial radius R_{vir} are the galaxy identifier and radial coordinate.

$$\rho_H(r) = \frac{M_{vir}}{4\pi R_{vir}^3}\,\frac{c^2\,g(c)}{x\,(1+cx)^2}\ , \qquad (3a)$$

where $x \equiv r/R_{vir}$ is the radial coordinate, c is the concentration parameter, and $g(c) = [\ln(1+c) - c/(1+c)]^{-1}$. The parameter c is found to be a weak function of the halo mass, given by $c \approx 14\,\left(M_{vir}/10^{11}\ \mathrm{M}_\odot\right)^{-0.13}$ (Bullock et al, 2001, Dutton, 2006, Gnedin 2006). This leads to

$$V^2_{\rm NFW}(r) = V^2_{vir}\frac{c}{g(c)}\frac{g(x)}{x}\ , \qquad (3b)$$

with $V_{vir} = V(R_{vir})$. It is interesting to note that in this scenario the present–day circular velocity, which also includes a baryonic component arranged in a disk, is predicted to be a *Universal* function of radius, tuned by few galaxy parameters (Mo, Mao & White 1998). However, it is well known that observations of spiral galaxies favor density concentrations lower than those predicted for CDM by Eq. (3a): DM halos detected around spirals do not show the NFW central cusp in favor of a core-like structure (Gentile et 2005; van den Bosch & Swaters 2001; Swaters et al. 2003; Weldrake et al. 2003; Simon et al. 2005; Donato et al, 2004; Gentile et al. 2007).

Therefore, the reconstruction of the mass distribution of DM halos from observations *in parallel* with that emerging from N-body simulations is required not only as a normal scientific routine, but also in view of a theory-vs-observations likely disagreement.

As an alternative to the simulation method, we will support the URC paradigm by means of a set of proper observational data and we will derive an analytical form for this curve, valid from the galaxy center out to its virial radius and characterized by the halo mass as the galaxy identifier. In detail, we extend/improve the URC_0 in PSS a) by adopting a different halo profile, proper to describe the halo distribution out to the virial radius, b) by using a number of RCs substantially more extended than those in PSS and c) by exploiting the relationship between the disk mass M_D, and the virial galaxy mass M_{vir}, recently obtained by Shankar et al. (2006). This will allow to build an "observational" Universal Curve, $V_{URC}(R; M_{vir})$, extended out to R_{vir} and having the virial mass as the galaxy identifier. This curve is the observational counterpart of the universal ΛCDM NFW N-body generated profile.

While pointing that the concept behind the Universal Rotation Curve may be valid also for galaxies of different Hubble Types (see Salucci and Persic, 1997), but a number of issues are still open and will be dealt elsewhere:

i) Sa galaxies amount, by number, to less than 10% of the whole spiral population, and are important objects in view of the dual nature of their stellar distribution. They show RC profiles with a clear systematics with luminosity (Rubin et al. , 1985), but, not unexpectedly, with some difference from those of the URC_0 (Noordmeer, 2007).

ii) Dwarf spirals with $V_{opt} < 50 km/s$ are not well studied and included in the URC yet, also because in these objects the RCs do not coincide with the circular velocity, being significant the complex asymmetric drift correction.

iii) The kinematical properties of spirals of very high stellar disk mass are not presently investigated with a suitably large sample.

iv) A possible additional URC physical parameter (e.g. the surface stellar density) to take care of the (small) variance of the RCs profiles that seems to be unaccounted by the luminosity.

Finally, let us remind that, in a flat cosmology with matter density parameter $\Omega_M = 0.27$ and Hubble constant $H_0 = 71$ km s^{-1} Mpc^{-1}, at the present time, the halo virial radius R_{vir}, i.e. the size of the virialized cosmological perturbation of mass M_{vir} scales with the latter as:

$$R_{vir} = 259\Big(\frac{M_{vir}}{10^{12}\ \mathrm{M}_\odot}\Big)^{1/3}\ \mathrm{kpc}\ . \qquad (4)$$



Figure 1. The core radius vs. virial mass relations for the SE sample and the present work (solid line).

(see e.g. Eke et al. 1996).

2 THE UNIVERSAL HALO VELOCITY PROFILE

We assume that the DM mass distribution is described by the Burkert (1995) profile

$$\rho(r) = \frac{\rho_0\, r_0^3}{(r + r_0)\,(r^2 + r_0^2)}\ , \tag{5a}$$

where r_0 is the core radius and ρ_0 the effective core density, in principle two independent parameters. Correspondingly, the total halo mass inside radius r is given by $M_H(r) = 4\,M_0\,\left[\ln\left(1+\frac{r}{r_0}\right) - \tan^{-1}\left(\frac{r}{r_0}\right) + \frac{1}{2}\,\ln\left(1+\frac{r^2}{r_0^2}\right)\right]$ with $M_0 = 1.6\,\rho_0\,r_0^3$, so that:

$$V^2_{URCH}(r) = 6.4\ G\ \frac{\rho_0 r_0^3}{r}\left\{ln\left(1+\frac{r}{r_0}\right) - \tan^{-1}\left(\frac{r}{r_0}\right) + \frac{1}{2}ln\left[1+\left(\frac{r}{r_0}\right)^2\right]\right\}\ . \tag{5b}$$

Inside R_l this profile is indistinguishable from the halo term (2b) in the URC$_0$ (Salucci & Burkert, 2000, Gentile et al 2004). At larger radii, the mass diverges only logarithmically with radius and converges to the NFW velocity profile, provided that $r_0 \ll R_{vir}$.

We fit the set of individual and coadded RCs of PSS with $V_{URC}(R; M_D, \rho_0, r_0)$ and derive the model parameters M_D, ρ_0, r_0 (see Salucci and Burkert, 2001):

$$\log \frac{\rho_0}{\text{g cm}^{-3}} = -23.515 - 0.964\,\left(\frac{M_D}{10^{11}\,\text{M}_\odot}\right)^{0.31} \tag{6a}$$

and

$$\rho_0 = 5\times 10^{-24} r_0^{-2/3} e^{-(r_0/27)^2}\,\text{g cm}^{-3}\ . \tag{6b}$$

Eqs. (1)-(2a)-(5b)-(6) define the URC out to R_l, $V_{URC}(R, M_D, r_0)$, from the "baryonic perspective". Let us notice that, as result of the RC mass modeling, r_0, differently from M_D and ρ_0, has quite large fitting uncertainties, viz. $\delta r_0/r_0 = 0.3 - 0.5$. Following our empirical approach, we do not extrapolate the URCH determined inside R_l out to $R_{vir} \gg R_l$. In that this will be uncertain besides of unknown validity. This quantity will be derived in the next section.

3 THE URC OUT TO THE VIRIAL RADIUS

We overcome the two main limitations of the URC$_0$, its problematic extrapolation between R_l and R_{vir} and the uncertainty in the estimate of the core radius, by determining the latter by means of a new *outer* observational quantity, the halo virial velocity $V_{vir} \equiv [GM_{vir}/R_{vir}(M_{vir})]^{1/2}$, related to the virial mass through Eq. (4). In detail, we obtain V_{vir} from the disk mass, suitably measured from inner kinematics through its relationship to the virial mass found by Shankar et al. (2006)

that is well represented (i.e. within 5%, the actual relation we use is shown in Fig. 2 and given in the Code indicated in the Discussion) by:

$$M_D = 2.3 \times 10^{10}\ M_\odot \frac{[M_{vir}/(3\ 10^{11}\ \mathrm{M}_\odot)]^{3.1}}{1+[M_{vir}/(3\ 10^{11}\mathrm{M}_\odot)]^{2.2}}\ . \tag{7}$$

This relationship is a consequence of the existence of i) the universal stellar mass function, (Bauldry et al 2004, Bell et al 2003) and of ii) the cosmological halo mass function as indicated by N-body ΛCDM simulations .[3] Let us notice that this relationship is obtained without assuming any halo density profile, so that it can be combined with the mass modelling of the inner kinematics.

Let us first derive $R_D(M_{vir})$, the disk scale-length as a function of the halo mass, by inserting Eq. (7) in the relationship:

$$\log \frac{R_D}{\mathrm{kpc}} = 0.633 + 0.379\ \log \frac{M_D}{10^{11}\ \mathrm{M}_\odot} + 0.069\ \left(\log \frac{M_D}{10^{11}\ \mathrm{M}_\odot}\right)^2 \tag{8}$$

obtained in PSS. We note that no result of this work is affected by the observational uncertainties on the relationship Eq. (8).

It is worth to compute the radial extrapolation needed to reach R_{vir} from $R_l = 6R_D$, a quantity that can also be associated with the baryonic collapse factor $F = R_{vir}/R_D$; we find $F \approx 90 - 15\ log\frac{M_{vir}}{10^{11}M_\odot}$, i.e. about 25 times the disk size $\sim 3R_D$.

Eq. (7) in combination with Eq. (4) allows us to add, to the PSS set of kinematical data leading to the URC$_0$, a new observational quantity: $V_{vir}(M_D) = GM_{vir}/R_{vir}$, relative to the virial radius. Then, we determine the core radius not from the inner kinematics, but as the value of r_0 for which the velocity model described by Eqs. (1), (2a), (5b) and (6a), matches (at R_{vir}) the virial velocity V_{vir} given by Eq. (7) and (4). Let us write this as:

$$\frac{GM_{vir}}{R_{vir}(M_{vir})} = V^2_{URCH}[R_{vir}(M_{vir}); \rho_0(M_{vir}), r_0)] \tag{9a}$$

where $\rho(M_{vir})$ as a short form for $\rho(M_D(M_{vir}))$ with Eq. (7) inserted in Eq. (6a). From the PSS inner kinematics we get the values of ρ_0 and M_D according to Eq. (6a), so that Eq. (9a) becomes an implicit relation between r_0 and M_{vir} (c_1, c_2 are known numerical constants):

$$M_{vir} = c_1 \frac{\rho_0(M_{vir}) r_0^3}{c_2 M_{vir}^{1/3}} \left\{ ln\left(1+\frac{c_2 M_{vir}^{1/3}}{r_0}\right) - \tan^{-1}\left(\frac{c_2 M_{vir}^{1/3}}{r_0}\right) + \frac{1}{2} ln\left[1+\left(\frac{c_2 M_{vir}^{1/3}}{r_0}\right)^2\right]\right\}\ . \tag{9b}$$

The above can be numerically solved for any M_{vir}, and the solution can be approximated by:

$$\log\left(r_0/\mathrm{kpc}\right) \simeq 0.66 + 0.58\ \log\left(M_{vir}/10^{11}\ \mathrm{M}_\odot\right)\ . \tag{10}$$

(an higher order approximation is given in the Code indicated in the Discussion). Let us stress that the present derivation of r_0 is very solid with respect to observational uncertainties: errors up to a factor 2 in M_{vir} in eq (7) trigger errors in r_0 lower than 40%, and errors in the outer halo velocity slope ($0.1 \leq R/R_{vir} \leq 1$) lower than 0.1. This is certainly smaller than the scatter of values with which this quantity is found by N-Body simulations and by SPH/semi analytical studies of galaxy formation including the baryonic components.

It is worth investigating a number of recently published super-extended (SE) RCs (Donato et al. 2005, Gentile et al 2004, Salucci et al 2003). They reach a radius larger than 5% (and up to 15%) of the virial radius, i.e. a radius at least twice as extended as those of the sinthetic curves in PSS. The mass modelling of these SE RCs (made in the original papers) shows an r_0 vs. M_{vir} relationship that is in good agreement with Eq. (10). Relation (10) and the above individual values differ by 20% -40% from those determined from the inner kinematics alone and given by means of eq (6b). Since in this paper (also because $R_l \sim r_0$), eq (6) is considered a prediction of the inner mass modelling rather than an actual measurement, such good agreement indicates the soundness of the PSS mass modelling.

Let us notice that only for a range of values of the crucial quantity $V(R_l)$- V_{vir}, with the first term obtained by the inner kinematics and the second one via Eqs. (4) and (7), there is a solution for eq.(9b), therefore, the existence of eq (10) and the agreement of the values of the Burkert core radii, measured independently at $0.05R_{vir}$, $0.1R_{vir}$ and R_{vir} are important tests passed by this profile.

Then, by means of Eqs. (1), (2a)-(5b)-(6a)-(10), we construct the full URC, extended out to the virial radius and with the virial mass as the galaxy indicator. It is useful to show the relationships we use (see Fig. 2). The mass model includes a Burkert DM halo of central density ρ_0, of core radius of size r_0 and a Freeman disk of mass M_D. The URC fits nicely the available velocity data out to R_l and it is valid out to the virial radius, where it exactly matches V_{vir}. Moreover, since M_{vir} is the quantity that in theoretical studies identifies a galaxy, we overcome the main limitation of URC$_0$.

We consider all of the three coordinate systems $r, r/R_D, r/R_{vir}$ equivalent to represent the main structural properties of

[3] There is no inconsistency in adopting the ΛCDM halo mass function and cored halo mass models, in that the latter can be formed astrophysical from the cosmological cuspy ones. Since both functions account the same cosmological objects, the Jacobian of their transformation defines a relation between the disk and virial mass in spirals (see Shankar et al. 2006).



Figure 2. The various relationships used in this paper.



Figure 3. The Universal Rotation Curve in physical units. Each curve corresponds to $M_{vir} = 10^{11}\ 10^{n/5} \mathrm{M}_\odot$, with $n = 1 \ldots 9$ from the lowest to the highest curve.

the mass distribution in spirals, but each of them showing some particular aspects. More specifically, it is then possible and useful to build several "URCs" i.e. $V_{URC}(r/R_{coo}; P)$, where P is a galaxy identifier (M_D, M_{vir}, L) and R_{coo} a radial coordinate $(r, r/R_D, r/R_{vir})$. Although not all these URCs are independent in a statistical sense, they all are relevant in that they all well reproduce the individual RCs and each of them highlights particular properties of the mass distribution.

In Fig. 3 we show $V_{URC}(r; M_{vir})$, the URC in physical units with the objects identified by the halo virial mass; each line refers to a given halo mass in the range $10^{11}\,\mathrm{M}_\odot \lesssim \mathrm{M_{vir}} \lesssim 10^{13}\,\mathrm{M}_\odot$; the halo mass determines both the amplitude and the shape of the curve. Note the contribution of the baryonic component, negligible for small masses but increasingly important in the larger structures, mirrors the behavior of the $M_{vir} - M_D$ relation. The general existence of an inner peak is evident but, especially at low masses, it is due to both dark and stellar components. Remarkably, the maximum value of the circular velocity occurs at about 15 ± 3 kpc, independent of the galaxy mass: this seems to be a main kinematic imprint of the DM - luminous mass interaction occurring in spirals. Furthermore, Fig. 3 shows that the "Cosmic Conspiracy" paradigm has no observational support: there is no fine tuning between the dark and the stellar structural parameters to produce the same particular RC profile in all objects (e.g. a flat one). Conversely, a number of relationships between the various structural parameters produce



Figure 4. The Universal Rotation Curve, normalized at its virial value $V_{URC}(R_{vir})$, as a function of normalized dark radius $x \equiv R/R_{vir}$. Each curve, from the highest to the lowest, corresponds to M_{vir} defined as in Fig. 3. The bold line is the NFW velocity profile (see text).

a variety of RC profiles. Moreover, the peak velocity of the stellar component $V_{disk}^{peak} = V_D(2.2R_D) = GM_D/R_D\ k$, with $k = const$, is not a constant fraction of the virial velocity as is found in ellipticals, (i.e $\sigma \propto V_{vir}$), but it ranges between the values 1 and 2 depending on the halo mass.

Moreover, as in the NFW (and Burkert) RC profiles, the URC profiles are found (moderately) decreasing over most of the halo radial extent. The paradigm of flat rotation curves is obviously incorrect even/especially intended as an asymptotic behavior at large radii. In fact, we find that both $V(0.05R_{vir})$,the velocity at the farthest radius with available kinematical in PSS and $V(3R_D)$, a main reference velocity of the luminous regions of spirals, are significantly $(10\% - 30\%)$ higher than the (observational) value of V_{vir}. This rules out a $V = constant$ extrapolation of the inner RCs out to regions non mapped by the kinematics and DM dominated regions. We note that this result is independent of the adopted halo density profile and is far from being granted on theoretical grounds.

In Fig. 4 we frame the URC from a full DM perspective by plotting $V_{URC}(R/R_{vir}; M_{vir})$. We set the virial mass M_{vir} as the galaxy identifier and R/R_{vir} as the radial "dark" coordinate, thus normalizing the amplitudes by $V_{vir} \propto M_{vir}^{1/3}$. This ensemble of curves, a main goal of the present work, is parallel to those emerging in N-body simulations and aims to represent the actual velocity profiles of spirals. In these variables the DM halos are self-similar; the whole system is self-similar in the outer regions, while in the innermost 30% of the halo size the baryons have influenced the dynamics and broken the self-similarity. In these coordinates it easily emerges that the maximum of the RC occurs at very different radii, viz. at $\simeq 2R_D$ for the most massive objects and at $\sim 10R_D$ for the least massive ones. Then, no reference circular velocities, to be considered as the actual physical counterparts of the empirical velocities of the Tully-Fisher relationship, exist in actual galaxy rotation curves.

In Fig. 5 we zoom into the URC to look for the inner (luminous) regions of spirals from a baryonic perspective: the URC is so expressed as a function of the "baryonic" radial coordinate r/R_D. This figure corresponds to Fig. 4 of PSS, with the important difference that here the virial mass, rather than the galaxy luminosity, is the galaxy identifier. Plainly, an inverse correlation between the average steepness of the RC slope and the halo mass holds, similar to the slope-luminosity relationship found by Persic & Salucci (1988). In this coordinate the stellar matter is closely self-similar, and the different shapes of the RC's curves are mainly due to the $M_{vir} - M_D$ relation. In the space defined by normalized circular velocity - dark radius - halo mass, spirals do not occupy random positions, but a well defined plane of very small thickness. We clearly see that, by filling only less than $10^{-3}\%$ of the available volume, the available kinematics of spiral galaxies defines the Universal Rotation Curve. Let us notice that, in principle, theories of the formation of spirals do not trivially imply the existence of such a surface that underlies the occurrence of a strong dark - luminous coupling.

We now show the URC DM density distribution. In Fig. 6 we show it as a function of x and M_{vir}. For $x < 10^{-1}$ the well known core-cusp discrepancy emerges, i.e. the DM density of actual halos around spirals is about one order of magnitude smaller and radially much more constant than the NFW predictions. At $x > 0.4$, for a concentration parameter $c = 13(M_{vir}/10^{12}\ M_\odot)^{-0.13}$, the observed halo densities are consistent with the NFW predictions for halos of the same virial mass. Note that this is a direct test: for halos with density profiles at $x > 0.5$ very different from the Burkert or the NFW profiles, Eq. 9b does not have solution.

More specifically, let us constrain the analytical form of the outer DM distribution. For $2 \leq r/r_0 \leq 18$, the following



Figure 5. The inner Universal Rotation Curve, normalized at its value at $4R_D$, as a function of normalized stellar radius R/R_D for galaxies with M_{vir} as in Fig. 3.



Figure 6. The URC halo density vs. the NFW halo density of objects of the same mass, as a function of normalized radius and virial mass. The axes labels are x, $logM_{vir}/M_{\odot}$ and $log(\rho/(g\ cm^{-3}))$

approximation for the Burkert and NFW profile holds ($y \equiv r/r_0$, $\epsilon = 0$)

$$V_{URCH}(y) = V_{URCH}(3.24)\frac{2.06\, y^{0.86}}{1.59 + y^{1.19+\epsilon}}\ . \tag{11}$$

Let us suppose that the actual *outer* DM velocity *profile* is different from the Burkert/NFW given by Eq. (11), i.e. $\epsilon \neq 0$. Then in Fig 7 we show that, even assuming large uncertainties in V_{vir}, in order to match both $V(R_l)$ and V_{vir}, we must have $\epsilon < 0.1$. This is a first direct support for the Burkert and the NFW density law to be able to represent the outer regions ($0.3 \leq r/R_{vir} \leq 1$) of DM galaxy halos.

Notice that weak-lensing shear fields, at several hundreds kpc from the galaxy centers, are found *compatible* with the predictions of the NFW density profile, but cannot exclude non-NFW profiles (Kleinheinrich ., et al., 2006 and references therein).

4 DISCUSSION AND CONCLUSIONS

In this paper, we have built the Universal Rotation Curve of spiral galaxies by means of kinematical and photometric data. We physically extended the URC, established for the inner region of galaxies in PSS out to R_{vir} and have been able to employ the virial mass M_{vir} as the parameter that characterizes spiral galaxies, and the virial radius R_{vir} as a unit of measure for the radial coordinate. This URC is meant to be the *observational* counterpart of the NFW rotation curve, emerging from cosmological simulations performed in the CDM scenario. The URC yields the gravitational potential at any radius and it allows to link the local properties in the inner luminous regions with the global properties of the DM halos.

DM halos have one (and likely just one) characteristic length scale, $r_0 \propto M_{vir}^{0.6}$, which it is not *naturally* present in current scenarios of galaxy formation. Thus, they do not show any sign of an inner cuspy region of size $r_s \propto M_{vir}^{0.4}$. The halo velocity contribution V_{URCH} rises with radius like a solid body at $r \sim 0$, decelerates to reach a maximum at 3.24 r_0 from where it start to slowly decrease out to R_{vir} with a slope that it is consistent with that of the NFW halos. The main significance of the URC concerns the full mass distribution (MD). First, it is possible to immediately exclude the following scenarios (and combinations of them): i) individual behavior, every object has its own MD; ii) unique behavior, every object has almost the same MD. Instead, the MD in spirals shows a remarkable *mass-dependent* systematics: both the dark and the stellar matter are distributed according to *profiles* that are functions of the total mass M_{vir} (see Fig. 8). Finally, the DM halo becomes the dominant mass component in galaxies at different radii, according to the galaxy mass: from $\sim 10^{-2} R_{vir}$ for the lowest masses, to $\sim 10^{-1} R_{vir}$ for the highest ones.

We write the equilibrium velocity of the halos around spirals as the following approximation of the relations in the previous Sect.:

$$V_{URCH} = A(M_{vir}) x^{-1/2} \Big\{ ln\Big[1 + \gamma(M_{vir})x\Big] - \tan^{-1}\Big[\gamma(M_{vir})x\Big] + \frac{1}{2} ln\Big[1 + \gamma(M_{vir})x^2\Big] \Big\}^{0.5}$$

with $A(M_{vir}) = 0.406 + 1.08\ log[M_{vir}/(10^{11}\mathrm{M}_\odot)] - 0.688\ \{\log[\mathrm{M_{vir}}/(10^{11}\mathrm{M}_\odot)]\}^2) + 0.766\ \{\log[\mathrm{M_{vir}}/(10^{11}\mathrm{M}_\odot)]\}^3$ and $\gamma(M_{vir}) = 26.78\ [M_{vir}/(10^{11}\mathrm{M}_\odot)]^{-0.246}$. This is the observational counterparts of N-body outcomes.

A Mathematica code for the figures in this paper is available at: http://www.novicosmo.org/salucci.asp.

5 APPENDIX

In this Appendix we discuss the observational evidence for the URC claim, the nature and the implications of which it is worth to clarify. The paradigm states that, when binned by luminosity, the RCs form a set of smooth, low-scatter synthetic curves, whose profiles and amplitudes are strong functions of the luminosity bin. [4]. Furthermore, the URC paradigm implies: i) rotation velocity slopes *vs.* rotation velocity amplitudes relationships (see fig (2) and (3) of PSS) and ii) a set of relations (Radial Tully Fisher relationship) holding at different radii x, defined as:

$$log\ V(x) = a_x M + b_x$$

where $x \equiv R/R_D$ and a_x and b_x are the fitting parameters, and M is the galaxy magnitude (Persic and Salucci 1991).

Evidence for the URC claim and/or its above implications comes from: a) detailed analyzes of independent samples: Catinella et al 2006, (2200 RC's, see their Fig. 12), Swaters 1999, (60 extended RC's, see Chapter 4); b) independent analyzes of the PSS sample: Rhee (1996), Roscoe (1999); c) the finding of a very tight RTF in PSS and other three different samples Willick (1997, see below), Yegorova et al. (2007).

The claim has been also tested by comparing the RCs of two samples of spirals (Courteau, 1998, 131 objects; Verheijen,

[4] The *analytical form* of the URC is built by assuming reasonable disk-halo velocity profiles, with three free parameters $(V(R_{opt}), a, \beta)$ that are obtained by χ^2-fitting the sinthetic curves.



Figure 7. Halo velocities at R_l and R_{vir} (*filled circles*) vs. the URC-halo velocity, given by Eqs. (11) (*solid line*) and vs. velocity profiles with average logarithmic slope steeper or shallower by an amount $\epsilon = 0.1$ (*dashed line*).



Figure 8. The dark halo (solid line), the disk (short dashed line), and the total (long dashed) URC mass profile for reference masses of $10^{11}M_\odot$ and $10^{13}M_\odot$. The distribution for the intermediate halo masses can be derived from section 2

1997, 30 objects) with the circular velocities predicted by the URC_0, once that the values of galaxy luminosity and disk length-scale are inserted in it. The face-value result of the test: 2/3 of the RCs are in pretty good agreement with the universal curve, while 1/3 show some disagreement, indicates that the URC is a useful tool to investigate the systematics of the mass distribution in spirals, but also it questions about its universality. However, while some of this disagreement may reflect an inefficiency of the URC_0 to reproduce the RCs, the actual performance of the URC is better than it is claimed. In fact, spurious data vs predictions disagreements are created in performing this test and precisely when they insert in URC_0 the